



# AAPC-CPC<sup>Q&As</sup>

Certified Professional Coder (CPC) Exam

**Pass AAPC AAPC-CPC Exam with 100% Guarantee**

Free Download Real Questions & Answers **PDF** and **VCE** file from:

<https://www.pass4itsure.com/aapc-cpc.html>

100% Passing Guarantee  
100% Money Back Assurance

Following Questions and Answers are all new published by AAPC  
Official Exam Center

- ⚙️ **Instant Download** After Purchase
- ⚙️ **100% Money Back** Guarantee
- ⚙️ **365 Days** Free Update
- ⚙️ **800,000+** Satisfied Customers



**QUESTION 1**

Which entity offers compliance program guidance to form the basis of a voluntary compliance program for a provider practice?

- A. Centers for Medicare and Medicaid Services (CMS)
- B. American Medical Association (AMA)
- C. Office of Inspector General (OIG)
- D. Office for Civil Rights (OCR)

Correct Answer: C

---

**QUESTION 2**

The mediastinum is:

- A. A location in the chest, bounded by the sternum, diaphragm, and lungs
- B. A small endocrine organ behind the heart
- C. A part of the lymphatic system
- D. Both the heart and lungs

Correct Answer: A

---

**QUESTION 3**

A 60-year-old male has three-vessel disease and supraventricular tachycardia which has been refractory to other management. He previously had pacemaker placement and stenting of LAD coronary artery stenosis, which has failed to solve the problem. He will undergo CABG with autologous saphenous vein and an extensive modified MAZE procedure to treat the tachycardia.

He is brought to the cardiac OR and placed in the supine position on the OR table. He is prepped and draped, and adequate endotracheal anesthesia is assured. A median sternotomy incision is made and cardiopulmonary bypass is initiated. The endoscope is used to harvest an adequate length of saphenous vein from his left leg. This is uneventful and bleeding is easily controlled. The vein graft is prepared and cut to the appropriate lengths for anastomosis. Two bypasses are performed: one to the circumflex and another to the obtuse marginal. The left internal mammary is then freed up and it is anastomosed to the ramus, the first diagonal, and the LAD. An extensive maze procedure is then performed and the patient is weaned from bypass. At this point, the sternum is closed with wires and the skin is reapproximated with staples. The patient tolerated the procedure without difficulty and was taken to the PACU.

Choose the procedure codes for this surgery.

- A. 33533, 33257, 33519, 33508
- B. 33535, 33259, 33519, 33508



C. 33533, 33257-51, 33519-51, 33508-51

D. 33535, 33259 51, 33519-51, 33508-51

Correct Answer: A

---

#### QUESTION 4

View MR 005398 MR 005398 Operative Report Preoperative Diagnosis: Nonfunctioning right kidney with ureteral stricture. Postoperative Diagnosis: Nonfunctioning right kidney with ureteral stricture. Procedure: Right nephrectomy with partial ureterectomy. Findings and Procedure: Under satisfactory general anesthesia, the patient was placed in the right flank position. Right flank and abdomen were prepared and draped out of the sterile field. Skin incision was made between the 11th and 12th

ribs laterally. The incision was carried down through the underlying subcutaneous tissues, muscles, and fascia. The right retroperitoneal space was entered. Using blunt and sharp dissection, the right kidney was freed circumferentially. The

right artery, vein, and ureter were identified. The ureter was dissected downward where it is completely obstructed in its distal extent. The ureter was clipped and divided distally. The right renal artery was then isolated and divided between 0

silk suture ligatures. The right renal vein was also ligated with suture ligatures and 0 silk ties. The right kidney and ureter were then submitted for pathologic evaluation. The operative field was inspected, and there was no residual bleeding

noted, and then it was carefully irrigated with sterile water. Wound closure was then undertaken using 0 Vicryl for the fascial layers, 0 Vicryl for the muscular layers, 2-0 chromic for subcutaneous tissue, and clips for the skin. A Penrose drain

was brought out through the dependent aspect of the incision. The patient lost minimal blood and tolerated the procedure well.

What CPT?coding is reported for this case?

A. 50234

B. 50220

C. 50230

D. 50240

Correct Answer: B

---

#### QUESTION 5

A business requires drug testing for cocaine and methamphetamines prior to hiring a job candidate. A single analysis with direct optical observation is performed, followed by a confirmation for cocaine. Which codes are used for reporting the testing and confirmation?

A. 80305 x 2, 80353

B. 80306 x 2, 80353



C. 80305, 80353

D. 80306, 80375

Correct Answer: C

[Latest AAPC-CPC Dumps](#)

[AAPC-CPC Practice Test](#)

[AAPC-CPC Exam  
Questions](#)