



MCQS^{Q&As}

Multiple-choice questions for general practitioner (GP) Doctor

Pass Test Prep MCQS Exam with 100% Guarantee

Free Download Real Questions & Answers **PDF** and **VCE** file from:

<https://www.pass4itsure.com/mcqs.html>

100% Passing Guarantee
100% Money Back Assurance

Following Questions and Answers are all new published by Test Prep
Official Exam Center

- ⚙️ **Instant Download** After Purchase
- ⚙️ **100% Money Back** Guarantee
- ⚙️ **365 Days** Free Update
- ⚙️ **800,000+** Satisfied Customers





QUESTION 1

33-year-old female complains of diplopia on upright gaze. Exam: ptosis can be seen. There are no other co significant Past medical history.

What is the most appropriate test for him?

- A. Ophthalmoscopy
- B. Checking red reflex
- C. Thyroid function test
- D. Visual field test

Correct Answer: B

This is a wrong key. Right key should be D. CT. A case of 3rd nerve palsy. So to cause CT should be done. Red reflex is seen to diagnose cataract.

QUESTION 2

Female with recently inserted IUCD coming with watery brownish vaginal discharge and abdominal pain what is the most likely diagnosis?

- A. Bacterial vaginosis
- B. D-ectopic pregnancy
- C. Uterine rupture
- D. Ovarian torsion

Correct Answer: A

QUESTION 3

At what CD4 count should antiretroviral treatment commence in asymptomatic HIV patients?

- A. Testosterone
- B. Growth hormone
- C. Somatostatin
- D. Oestrogen
- E. Somatostatin



Correct Answer: D

The growth spurt at puberty is brought about by the secretion of androgens in the male and oestrogens in the female. However, it is oestrogens that ultimately terminate growth by causing the epiphyses in the long bones to fuse. Thus oestrogens rather than androgens are responsible for skeletal maturation, epiphyseal fusion and cessation of growth in males and females. All the other hormones are involved in growth alone

QUESTION 4

A terminally ill with metastatic carcinoma presents with dysphagia and difficulty in swallowing.

What is possible treatment?

- A. Analgesic
- B. Cotrimazole
- C. PO fluconazole
- D. Nystatin suspension
- E. Amphotericin B IV

Correct Answer: C

PO fluconazole. Treatment of carcinoma can predispose to development of esophageal candidiasis as follows: Oral fluconazole (200 ?400mg) daily for 14 to 21 days (if needed IV fluconazole can also be given).

QUESTION 5

Which of the following will have the lowest glucose level on pleural effusion?

- A. Cancer
- B. Pulmonary tuberculosis
- C. Rheumatoid arthritis
- D. Pneumonia

Correct Answer: C

Rheumatoid arthritis has the lowest glucose level of all the causes of pleural effusion described here.

[Latest MCQS Dumps](#)

[MCQS VCE Dumps](#)

[MCQS Exam Questions](#)